

BEST AVAILABLE COPY

LETTERS PATENT

Number 514732

ELIZABETH THE SECOND, by the Grace of God Queen of New Zealand and Her Other Realms and Territories, Head of the Commonwealth, Defender of the Faith; To all to whom these presents shall come, Greeting:

WHEREAS pursuant to the Patents Act 1953 an application has been made for a patent of an invention for

Method for screening crystallization conditions in solution crystal growth

(more particularly described in the complete specification relating to the application)

AND WHEREAS

THE UAB RESEARCH FOUNDATION, 1120 G Administration Building, Birmingham, Alabama 35294, United States of America

(hereinafter together with his or their successors and assigns or any of them called "the patentee") is entitled to be registered as the proprietor of the patent hereinafter granted:

Address for service: A J PARK, 6th Floor, Huddart Parker Building, 1 Post Office Square, Wellington, New Zealand


NOW, THEREFORE, We by these letters patent give and grant to the patentee our special licence, full power, sole privilege, and authority, that the patentee by himself, his agents, or licensees and no others, may subject to the provisions of any statute or regulation for the time being in force make, use, exercise and vend the said invention within New Zealand and its dependencies during a term of twenty years from 5 April 2000 and that the patentee shall have and enjoy the whole profit and advantage from time to time accruing by reason of the said invention during the said term:

AND WE strictly command all our subjects whomsoever within New Zealand and its dependencies that they do not at any time during said term either directly or indirectly make use of or put into practice the said invention, nor in any way imitate the said invention without the consent, licence, or agreement of the patentee in writing under his hand, on pain of incurring such penalties as are prescribed by law and of being answerable to the patentee according to law for his damages thereby occasioned:

PROVIDED ALWAYS:

- (1) That these letters patent shall determine and become void if the patentee does not from time to time pay the renewal fees prescribed by law in respect of the patent:
- (2) That these letters patent are revocable on any of the grounds prescribed by the Patents Act 1953 as grounds for revoking letters patent:
- (3) That nothing in these letters patent shall prevent the granting of licences in the manner in which and for the considerations on which they may by law be granted:
- (4) That these letters patent shall be construed in the most beneficial sense for the advantage of the patentee.

IN WITNESS whereof We have caused these letters patent to be signed and sealed on 10 May 2004 with effect from 5 April 2000.



Neville Harris
Commissioner of Patents, Trade Marks and Designs

**METHOD FOR SCREENING CRYSTALLIZATION
CONDITIONS IN SOLUTION CRYSTAL GROWTH**

Field of the Invention

The present invention relates to the crystallization
5 of proteins in or from protein solutions. The present
invention particularly relates to a method of screening a
large number of protein crystal growth conditions which
may be conducive to protein crystallization. Even more
particularly, the present invention relates to a method
10 which identifies one or more optimum protein crystal
growth conditions, while at the same time using
substantially less protein solution.

Background of the Invention

The crystallization of proteins for structure-
15 function studies and structure based drug design has
become an increasingly important part of biotechnology
research. When crystal growth is attempted for a new
protein, the appropriate chemical conditions (i.e.
protein concentration in solution, precipitate type and
20 concentration, pH, and growth temperature) are unknown
and have typically been determined by trial and error
experimentation.

Typically 1000 or more different sets of crystal
growth conditions are screened to determine conditions
25 conducive to crystallization. The screening involves
repetitive procedures that are extremely laborious and
tedious. With present laboratory protein crystal growth
equipment, each crystallization chamber requires about
one micro-liter of protein solution. The protein
30 solutions typically have concentrations in the range of
10 to 25 micrograms per microliter to facilitate crystal
growth. Therefore, to screen 1000 samples typically

(followed by page 2a)

requires between 10 and 25 milligrams of protein. This is a considerable and costly amount, especially for proteins that are difficult to isolate or generally express. A large percentage (about 50%) of the proteins
5 cannot easily be expressed in milligram quantities.

Thus, it would be desirable to provide methods for screening protein crystal growth conditions that require picogram to microgram amounts of protein for each screening condition. Preferably such methods would
10 require only picogram to nanogram amounts of protein in picoliter to nanoliter volumes in each screening condition sample.

It would be further desirable to provide high throughput screening methods for screening protein
15 crystal growth conditions in a large number of samples on a sub-microgram scale. These methods would use a microarray as a platform for protein crystal growth. The methods would also utilize automatic dispensing of solutions and scoring of crystal growth.

20 It is an object of the present invention to fulfil these needs or at least to provide the public with a useful choice.

Summary of the Invention

The present invention concerns a method of screening
25 protein crystal growth conditions employing a minimal amount of protein, on a picogram to microgram scale. Each screening sample has picogram to microgram amounts of protein in a picoliter to nanoliter volume. Predetermined protein crystal growth conditions are
30 maintained and crystal growth is analyzed using both qualitative and quantitative criteria.

INTELLECTUAL PROPERTY
OFFICE OF N.Z.

21 NOV 2003

(followed by page 2b)

More specifically, in one aspect the present invention provides a method of screening protein crystal growth conditions employing picogram, nanogram or microgram amounts of protein comprising the steps of:

providing a microarray with a plurality of wells for protein solution and a plurality of wells for precipitate solution, wherein said microarray has micro-channels for fluid communication between said wells for protein solution and said wells for precipitate solution;

accurately dispensing a volume of from said about 0.001 nl to about 250 nl of a protein solution into said wells for protein solution;

controlling the protein crystal growth conditions of each of said wells so that the protein crystal growth conditions in at least two of said wells differ, wherein controlling crystal growth includes adding precipitate solution to the wells for precipitate solution, and wherein protein solution and precipitate solution are in fluid communication via said micro-channels;

affixing a cover to the microarray such that said wells form micro-chambers and said micro-channels form capillary tubes, said cover providing an optically clear path from said micro-chambers; and

observing protein crystal growth or protein precipitation in said wells.

In a further aspect, the present invention provides a microarray for screening protein crystal growth at nanogram or picogram protein amounts comprising:

a plurality of wells wherein said wells are adapted for holding volumes of protein solution from about 0.001 nl to about 250 nl;

(followed by page 3)

a separate plurality of wells for holding a precipitate solution, and wherein said microarray has micro-channels for fluid communication between wells for holding protein solution and wells for holding precipitate solution;

and further wherein the material of the microarray comprising said wells is minimally water absorbing; and an optically clear path from said wells.

In another embodiment the present invention provides a microarray for screening protein crystal growth at nanogram or picogram protein amounts comprising a plurality of wells wherein said wells are adapted for holding volumes of protein solution from about 0.001 nl to about 250 nl and are in fluid communication through micro-channels with another plurality of wells for holding precipitate solution.

In a preferred embodiment, a microarray is provided for use in methods of screening protein crystal growth. Preferably the microarray has a plurality of micro-chambers in the microarray. The micro-chambers may be

passive or a combination of passive micro-chambers that are connected with miniaturized active control elements such as, but not limited to, valves, pumps and electrodes. A protein solution is automatically
5 dispensed into the micro-chambers. The protein crystal growth conditions of each of the micro-chambers is adjusted so that the protein crystal growth conditions of at least two of the micro-chambers differs. Protein crystal growth in the micro-chambers is then analyzed
10 based on both the qualitative amount of crystallization and the quality of the crystals formed.

Additional objects, advantages, and features of the present invention will become apparent from the following description and appended claims, taken in conjunction
15 with the accompanying drawings.

Brief Description Of The Drawings

The various advantages of the present invention will become apparent to one skilled in the art by reading the following specification and subjoined claims and by
20 referencing the following drawings in which:

Figure 1 is a schematic illustrating a two well design in a microarray;

Figure 2A is a schematic showing a top view of the placement of protein and precipitate solutions in a one
25 well design;

Figure 2B is a schematic showing a side view of placement of protein and precipitate solutions in a one well design;

Figure 2C is a schematic showing a side view of an
30 alternative placement of protein and precipitate solutions in one well;

Figure 2D is a schematic showing a side view of placement of protein and precipitate solutions in two wells;

Figure 2E is a schematic showing a top view of the placement of protein and precipitate solutions in a two well design;

Figure 3 is a photograph showing a microarray; and

Figure 4 is a photograph of a protein crystal obtained with nanogram amounts of protein in nanoliter volumes.

Detailed Description of Invention

The method of the present invention is for screening protein crystal growth conditions in protein solutions employing a minimal amount of protein in a minimal volume, on a pico, nano or meso scale. Pico, nano or meso scale as used herein preferably employs (on average) picogram (pg), nanogram (ng) or microgram (μg) amounts of protein in picoliter (pl) or nanoliter (nl) volumes. Preferably, the amount of protein in each screening sample is less than about 5 μg . More preferably, the amount of protein in a screening sample will be less than about 1 μg . In one embodiment, the volume of protein solution in a screening sample is preferably from about 0.001 nl to about 250 nl and more preferably about 0.01 nl to about 10 nl. It will be appreciated by those skilled in the art that the volumes actually employed for any particular protein will be a function of (without limitation) the target protein and its concentration in the protein solution.

The protein solution contains one or more desired proteins for crystallization. As used herein, the term "protein" is meant to include all naturally occurring and

synthetic peptides, polypeptides, proteins, and protein complexes. In one preferred embodiment the concentration of protein in the solution is from about 0.1 $\mu\text{g}/\mu\text{l}$ to about 50 $\mu\text{g}/\mu\text{l}$, more preferably from about 0.1 $\mu\text{g}/\mu\text{l}$ to about 10 $\mu\text{g}/\mu\text{l}$, and still more preferably about 0.1 $\mu\text{g}/\mu\text{l}$ to about 1.0 $\mu\text{g}/\mu\text{l}$. In another preferred embodiment, the solution is buffered to a pH between about 2.0 and about 10.0, more preferably from about 3.0 to about 8.5. If desired, the protein solution may optionally contain agents that aid in solubilizing the protein at the desired protein concentration or conditions. For example, if the protein is a membrane protein, the protein solution may optionally contain one or more surface active agents, such as a detergent mixture. In one preferred embodiment, the protein solution also comprises components that assist in crystallization. By way of non-limiting example, the protein solution will comprise an aqueous salt solution, polyethylene glycol, or an alcohol. Such components as well as their selection, ranges, contraindications and the like are well known to those skilled in the art. See, for example, Gilliland, G.L. et al., *Acta Crystallogr.* D50:408-413 (1994); McPherson, A., *Crystallization of Biological Molecules*, Cold Spring Harbor Press, Cold Spring Harbor, NY, pp. 487-524 (1999), expressly incorporated by reference.

The protein solution is dispensed onto a platform. The platform can be, by way of non-limiting example, a glass slide, a multi-well plate or a microarray. The solution is preferably dispensed using a device with picoliter or nanoliter accuracy. Preferably the dispensing device has at least a 90% accuracy on a

picoliter or nanoliter scale. The protein solution can be dispensed manually using, for example, a syringe. In a highly preferred embodiment, automatic dispensing devices are used to dispense the protein solution.

5 A second solution, the reservoir or precipitate solution is provided. The precipitate solution is a solution that helps to bring about protein crystallization. It can comprise, for example, a salt
10 second solution is provided either before, after, or simultaneously with the protein solution. The volume of the precipitate solution is typically equal to or greater than the volume of protein solution. The placement of the second solution is dependent on the crystallization
15 method used but is typically in fluid communication with the first solution. Fluid communication can be liquid-liquid, liquid-vapor or vapor-vapor communication. Generally, a channel is provided for fluid communication. A channel is broadly defined herein as a space that
20 enables fluid communication to occur. In the liquid-liquid diffusion method, the protein solution and precipitate solution contact each other at an interface. In batch crystallization, the two solutions are mixed together. If vapor diffusion crystallization is desired,
25 the two solutions are kept separate but space is allowed for the diffusion of vapor between the solutions. Or, in an alternate embodiment, a single source or reservoir of the second solution may be employed. In yet another alternate embodiment, a desiccant source or a dry gaseous
30 reservoir may be employed in place of the second solution. Specific conditions and variations in these methods are well known to the skilled artisan.

Protein crystal growth is monitored periodically, either qualitatively, quantitatively, or both. This may be by manual inspection using high resolution microscopy or electron microscopy. Preferably, protein crystal growth may be monitored automatically, by, for example, high resolution optical means which automatically detects crystal growth based on, for example, edge analysis. If desirable crystal growth is observed in a sample, the protein crystal growth conditions of that sample can be reproduced on a macro scale to produce a protein crystal for further analysis. Alternatively, if a precipitate or a clear sample is observed, these conditions can be used to optimize the conditions for additional screening. It will be appreciated that the platform must employ at least one path that is visually and/or optically clear to the method of detection.

In at least one preferred embodiment the method of the present invention for screening protein crystal growth employs a microarray with a plurality of wells or reservoirs as the platform. A microarray may be constructed, for example, similar to a micro-electromechanical chip. The microarray preferably has a planar shape and employs a size and thickness that are compatible with manual or automated plate grippers. The microarray can be made from different materials and by different techniques known to those skilled in the art. The material of the microarray that includes the wells or reservoirs is preferably minimally water absorbing, and is otherwise sufficiently unreactive with the components of the solution. This may be done as a laminate or provided as a coating, for example. Alternatively, a material that absorbs water at a predictable rate can

also be used to construct the wells or reservoirs. The volumes of protein and precipitate solutions may then be adjusted to compensate for the water absorption of the material. Preferred materials include, but are not
5 limited to, glass, fused silicon, quartz, a silicon wafer, a polymer or a polysulphone. Alternatively, the microarray can be made from a material coated with a hydrophobic material, such as polysulphone, to limit water absorption in the microarray. Alternatively, the
10 microarray comprises more than one material. Preferably, the microarray is a composite with a bottom of thin glass plate bonded to plastic, glass, silicon rubber or other materials in which wells can be manufactured, with at least one side providing an optical path that is
15 acceptable to the detection technique employed.

In an alternate embodiment, the surfaces of the wells are hydrophobic. For example, the material of the microarray may have a hydrophobic surface. Alternatively, the surfaces of the wells may be coated
20 with a hydrophobic coating. Although not necessary, the hydrophobic surfaces of the wells prevent the drops of solutions from spreading.

The microarray includes a multitude of micron sized wells on the surface of the chip. The term wells
25 encompasses wells, micro-chambers and any indentation sufficient of holding or retaining a desired volume of from about 0.001 nl to about 500 nl, preferably from about 0.01 nl to about 20 nl. The wells are spaced from each other on the surface. The precise number of wells on
30 the surface of the microarray can vary, and the total number of wells on the surface is a matter of choice for the user.

Each of the wells has a volume sufficient to hold an amount of protein solution adequate for growing a protein crystal. Preferably, each of the wells holds a volume from about 0.001 nl to about 250 nl, preferably from about 0.01 nl to about 10 nl.

The wells of the microarray are made by using an etchant such as hydrogen fluoride or by other known etching or fabrication techniques.

The wells can include known means for controlling conditions, individually or collectively, such as pressure, heating or cooling the wells, humidity levels in the wells as well as known means for stirring materials loaded into the wells.

In one arrangement, the wells of the microarray are not connected and separate from each other. In an alternative arrangement, adjacent wells of the microarray are connected by one or more channels which provide fluid communication between the adjacent wells (Figures 1 and 2D-E). Preferably, the connecting channels will have cross-section dimensions and length allowing control over the rate of transport of fluid, vapor, buffer, or precipitating or crystallizing agents through the channels. In one embodiment, varying the dimensions of the channels controls protein crystal growth condition. In an alternate embodiment, protein crystal growth conditions are controlled by placing a material in the micro-channels that controls fluid communication between the wells. Non-limiting examples are membranes, acrylamide or agarose. For example, the connecting micro-channels are from about 0.0001 to about 0.2 microns wide and from about 0.00005 to about 0.1 microns deep. Alternatively, the micro-channels are from about 0.0001

to about 2.0 microns wide and from about 0.00005 to about 0.5 microns deep. The micro-channels are formed in the microarray chip by the known etching techniques.

5 An example of two wells in a microarray (10) connected by a micro-channel is shown in Figure 1. The protein solution well 12 is connected to precipitate solution well 14 by a microchannel 16. The dimensions of each well are designed to hold the desired amount of solution and may have the same or different dimensions.
10 Initially, protein sample is dispensed into well 12 to an initial liquid height 18 and precipitate solution is dispensed into well 14 with liquid height 20. The top of the wells and microchannel are sealed by an optically clear cover 22. In vapor diffusion crystallization, the
15 precipitate solution in well 14 has a lower vapor pressure than the protein solution in well 12, causing diffusion of solvent from well 12 to well 14 until the solution liquid height in well 12 reaches a final height 24. The concentration of the protein solution in well 12
20 precipitates protein crystal formation.

The microarray can also include a known means for transmitting a fluid or gas to the wells of the microarray from an external source. For example, an external mechanical pumping system marketed by Watson-
25 Marlowe, Inc., under the trade designation "205U" can be used. The pumping system is a multi-channel cassette which delivers fluid or gas in reproducible and accurately controlled amounts.

Optionally, micro-valves are disposed in the wells
30 and micro-channels to regulate the flow of fluid or vapor between the wells and through the micro-channels in a known manner.

An automated dispensing mechanism capable of accurately and/or repeatedly dispensing picoliter and/or nanoliter volumes is also provided. Preferably, the automated dispensing mechanism has an accuracy of at least about 90%. The automated dispensing mechanisms are preferably Piezo-based or fast solenoid dispensing mechanisms. More preferable, the dispensing mechanism is a fast solenoid dispensing mechanism. The dispenser has a large number of parallel capillaries. The capillaries are in fluid communication with a source of protein solution, a source of precipitate solution, and a source of buffer solution. The dispensing can be actuated by ultrasonic transducers that efficiently produce a pressure wave in the capillaries that contain the solutions. The dispenser is analogous to ink jet printer heads for computer printers but the fluid is not heated, thus not damaging the solutions.

The protein solution preferably comprises an aqueous protein solution at a concentration of from about 0.1 $\mu\text{g}/\mu\text{l}$ to about 50 $\mu\text{g}/\mu\text{l}$. Preferably, the concentration is from about 0.1 $\mu\text{g}/\mu\text{l}$ to about 10 $\mu\text{g}/\mu\text{l}$, more preferably from about 0.1 $\mu\text{g}/\mu\text{l}$ to about 1.0 $\mu\text{g}/\mu\text{l}$. Preferably, the protein solution comprises a detergent mixture when crystallizing membrane proteins. The precipitate solution preferably comprises a concentrated aqueous salt solution or polyethylene glycol as precipitating agents. The buffer solution preferably has pH between about 2 and about 10.

The automated dispensing mechanism dispenses an initial volume of protein solution, an initial volume of precipitate solution, and an initial volume of buffer solution from the source of protein solution, the source

of precipitate solution, and the source of buffer solution; respectively, into preselected wells or connecting channels of the microarray.

5 The placement of the initial volume of protein solution, the initial volume of precipitate solution, and the initial volume of buffer solution in the preselected wells or channels of the microarray is dependent upon the method utilized to effect crystallization of the protein in the protein solution.

10 Preferred methods to effect crystallization of the protein in the protein solution include liquid-liquid diffusion, batch diffusion, and vapor diffusion.

15 In the liquid-liquid diffusion method, the initial volume of protein solution is placed in one set of preselected wells, and the initial volume of precipitate solution is placed in a separate or different set of wells. The protein solution wells are connected to the precipitate solution wells by micro-channels. The initial volume of buffer solution may be placed in the
20 micro-channels, or alternatively added directly to the initial volume of protein solution and/or precipitate solution.

The concentration, amounts, precipitate type, and pH of the initial volumes of protein solution, precipitate
25 solution, and buffer solution are primary conditions which determine protein crystal growth in a protein solution. In preparing the initial solutions, and in the automated dispensing mechanism placement, these conditions and the sample placement are varied in
30 accordance with a pre-designed program.

A cover plate is affixed to the microarray to convert the wells to micro-chambers and to convert the

micro-channels to a capillary tube structure. The cover plate can be made of the same or different material as the microarray, but the cover plate (or some portion of the well or chamber) must be transparent to permit optical analysis of the protein solutions in the chambers of the microarray. Preferably, the cover plate will be glass or other material that is visually or optically clear, such as an optically clear tape.

Alternatively, the environment surrounding the microarray can be controlled to limit evaporation of the solutions. Under controlled conditions of, for example, temperature and humidity, covering the samples may not be necessary.

The crystallizing agent in the precipitate solution, in selected micro-chambers, diffuses via the connecting capillaries to selected micro-chambers containing protein solution.

Protein crystal growth in the different chambers are then monitored by high resolution or other optical means which automatically detects crystal growth based on well known edge analysis. Alternatively, the protein crystal growth can be monitored by manual inspection using high resolution microscopy or electron microscopy. Preferably the protein crystal growth in the chambers is monitored by high resolution optical means which automatically detects crystal growth based on edge analysis.

Once crystal growth in a chamber is detected, that chamber's protein crystal growth conditions can be reproduced on a macro scale to produce a protein crystal which can be analyzed by x-ray crystallography. Alternatively, if a precipitate or clear sample is

observed, the conditions in those samples can be used to optimize conditions for additional screening.

In the vapor diffusion method, the initial volume of protein solution is placed in one set of preselected wells, and the initial volume of precipitate solution is placed in a separate or different set of wells based on a pre-designed program, as with the liquid-liquid diffusion method (Figures 2D-E). The protein solution wells are connected to the precipitate solution wells by micro-channels. The initial volume of buffer solution is added to the initial volume of protein solution and/or initial volume of precipitate solution. Alternatively, the protein solution and precipitate solution can be placed in the same well such that the two solutions do not come into contact (Figures 2A-C).

As with liquid-liquid diffusion, the crystal growth is varied in different wells in accordance with a pre-designed program in which the placement, concentration, amounts, precipitate type, and pH conditions are varied in the different wells.

A cover plate is then affixed to the microarray as with the liquid-liquid diffusion method. The vapor pressure of the precipitate solution is lower than the vapor pressure of the protein solution. This causes the protein solution in a micro-chamber which is connected via a capillary to a micro-chamber containing a precipitate solution to evaporate and become supersaturated causing precipitation of protein. Crystal growth is monitored as in the liquid-liquid diffusion.

Alternatively, the protein solution is placed into wells of the microarray and the microarray is exposed to a single reservoir with the precipitate solution. This

method allows for less fluid dispensing, but also less control of the protein crystal growth conditions with respect to each protein sample.

5 In the batch method, the volume of protein solution, the volume of precipitate solution, and the volume of buffer solution are placed together in individual wells of the microarray. In this method, the chip does not have connecting channels between the wells.

10 As with liquid-liquid diffusion and vapor diffusion methods, the crystal growth is varied in different wells in accordance with a pre-designed program in which the placement, concentration, amounts, precipitate type, and pH conditions are varied in the different wells.

15 As with liquid-liquid diffusion and vapor diffusion methods, a cover plate is affixed to the microarray, and the crystal growth is then monitored.

20 If desired, fluid or gas can be delivered to the micro-chambers in reproducible and accurately controlled amounts from an external source by the external mechanical pumping system described above. Gas can also be delivered from the pressure generated by a standard glass bottle or tank. The fluid or gas delivered to the micro-chambers can be regulated by the micro-valves. The fluid or gas can be used to further alter the crystal growth conditions of the micro-chamber and increase the size of the protein crystals grown. These protein crystals can then be harvested and examined by x-ray crystallography or nuclear magnetic spectroscopy or other appropriate techniques.

30 Advantages of the present invention should now be apparent. The present invention provides a method of screening protein crystal growth conditions on a nano or

meso scale. The method provides a means of screening protein crystal growth conditions for proteins that cannot be expressed in milligram quantities as well as those that can be expressed in larger quantities. Moreover, the substantial reduction in protein needed for the present invention reduces the costs associated with screening protein crystal growth conditions.

Disclosed is an apparatus for screening crystal growth conditions. The apparatus comprises a microarray for the protein and precipitate solutions, an automatic dispensing mechanism for dispensing the solutions and an automated means for analyzing crystal growth.

The desired solutions, i.e., protein, precipitate and a buffer, are preferably automatically dispensed at a preset picoliter or nanoliter volume into the microarray by an automated dispensing mechanism. Preferably, the automatic dispensing mechanism dispenses discrete drops. Screening conditions such as the type of buffer and pH can be varied from sample to sample by programming the automatic dispenser. For example, arbitrary screens varying pH could be programmed by mixing the proper ratios using different drop counts from different stock solutions having different pH values. A pH range from 2.0 to 10.0 is then screened in steps of 0.2-0.5 pH units. Other conditions, such as crystallization agents and salt concentration are also controlled in a similar manner.

Mixing of the reagents can either be done before dispensing or after the solutions are dispensed into the microarray. Mixing in the microarray, for example, can be accomplished by ultrasonic mixing, high-speed

dispensing of picoliter drops, rapid temperature fluctuation, or by static diffusion.

After mixing, preferably the wells of the microarray are sealed to control the microenvironment in the wells and to prevent evaporation of the solutions to dryness. More preferably, the wells are sealed with optically clear tape. Sealing the microarray involves an arm mounted on a YZ transverse mechanism. The X direction is along the plate transport direction. The arm, holding a roll of clear tape, moves past the last well in the row, drops to form positive vertical (Z axis) pressure, and then begins to move back in the negative Y direction while at the same time rotating the tape roll. Once the tape leaves the plate area, a guillotine mechanism shears the tape. The plate then moves in the X direction onto the next indexed row and the dispense process initializes again. Automated taping is reliably performed in many industries.

Protein crystal growth in the different wells is monitored by high resolution optical means which automatically detects crystal growth based on well known edge analysis. Such image acquisitions systems are commercially available.

The foregoing and other aspects of the invention may be better understood in connection with the following example, which is presented for purpose of illustration and not by way of limitation.

EXAMPLE 1

Nanoliter protein droplets were used for vapor diffusion, batch and liquid diffusion crystallization screening. The protein solutions of either lysozyme, thaumatin, or NAD synthetase were applied using a five

microliter Hamilton syringe. To ensure complete wetting of the small droplet to the experiment chamber, the tip of the Hamilton syringe was placed in contact with the wall of each experiment chamber. A variety of
5 microarrays were designed to accommodate protein solution droplets with volume ranges of 5-20 nanoliters and precipitate volumes of 100-200 nanoliters. The array prototyping was accomplished using MicroScope slides with permanent sealing of neoprene gaskets of varying
10 thickness (0.1 mm 0.5 mm). Once all solutions were applied to an individual experiment chamber within the microscope slide, the experiment was sealed (with oil or grease) by placing a glass cover slide over the top of the gasket. Figure 3 is a photograph of a typical design
15 for a 60 chamber array prototype (gasket thickness = 0.1 mm) and Figure 4 is a photograph of crystals that were grown to 10 nanoliter protein droplets using a similar microarray slide.

A Cartesian robotic dispensing system was used to
20 prepare crystallization solutions in a 6 by 10 experiment array. Five nanoliters of protein plus five nanoliters of precipitant were dispensed into one merged droplet in one depression in the experiment chamber (Figure 3) and 50 nanoliters of precipitant plus 50 nanoliters of buffer
25 were merged into one droplet in the connected depression. Thus, four solutions were dispensed for each experiment, and $6 \times 10 \times 4 = 240$ total for the entire 6 by 10 array. Cartesian's instrument was able to dispense all of the solutions in less than 20 minutes. All external
30 conditions used were known crystallization conditions for the particular proteins tested. The experiment was manually sealed and incubated at 22°C. for a period of

one day. Crystals were observed in seventy percent of the droplets. While not wishing to be bound by theory, it is believed that the failure to observe crystals in 30% of the wells was due to inaccurate dispensing of the protein and precipitant five nanoliter drops in that the peizo tip did not position the drops together.

From the above description of the invention, those skilled in the art will perceive improvements, changes and modifications. Such improvements, changes and modifications within the skill of the art are intended to be covered by the appended claims.

WHAT WE CLAIM IS:

1. A method of screening protein crystal growth conditions employing picogram, nanogram or microgram amounts of protein comprising the steps of:

providing a microarray with a plurality of wells for protein solution and a plurality of wells for precipitate solution, wherein said microarray has micro-channels for fluid communication between said wells for protein solution and said wells for precipitate solution;

accurately dispensing a volume of from said about 0.001 nl to about 250 nl of a protein solution into said wells for protein solution;

controlling the protein crystal growth conditions of each of said wells so that the protein crystal growth conditions in at least two of said wells differ, wherein controlling crystal growth includes adding precipitate solution to the wells for precipitate solution, and wherein protein solution and precipitate solution are in fluid communication via said micro-channels;

affixing a cover to the microarray such that said wells form micro-chambers and said micro-channels form capillary tubes, said cover providing an optically clear path from said micro-chambers; and

observing protein crystal growth or protein precipitation in said wells.

2. The method of Claim 1 wherein said protein solution comprises a component selected from the group consisting of buffers, surface active agents, salts, alcohol, polyethylene glycol and mixtures thereof.

3. The method of Claim 1 wherein said protein solution is buffered.

4. The method of Claim 1 wherein controlling protein crystal growth further comprises employing varying dimensions of the channels.

5. The method of Claim 1 wherein fluid communication is by liquid-liquid diffusion of said precipitate solution and said protein solution.

6. The method of Claim 1 wherein said precipitate solution has lower vapor pressure than the protein solution and fluid communication is by vapor diffusion.

7. The method of Claim 1 wherein the precipitate solution has a volume from about 0.001 nl to about 250 nl.

8. The method of Claim 1 wherein protein crystal growth or protein precipitation is observed by microscopy.

9. The method of Claim 8 wherein the microscopy is differential interference contrast microscopy.

10. The method of Claim 1 wherein the protein solution is dispensed into said wells by fast solenoid dispensing.

11. A microarray for screening protein crystal growth at nanogram or picogram protein amounts comprising:

a plurality of wells wherein said wells are adapted for holding volumes of protein solution from about 0.001 nl to about 250 nl;

a separate plurality of wells for holding a precipitate solution, and wherein said microarray has micro-channels for fluid communication between wells for

holding protein solution and wells for holding precipitate solution;

and further wherein the material of the microarray comprising said wells is minimally water absorbing; and
an optically clear path from said wells.

12. The microarray of Claim 11 wherein said wells further comprise a material that is substantially hydrophobic.

13. The microarray of Claim 11 wherein at least two micro-channels have different dimensions.

14. The microarray of Claim 11 wherein said microarray further comprises a means for sealing to prevent evaporation from said wells.

15. A microarray for screening protein crystal growth at nanogram or picogram protein amounts comprising a plurality of wells wherein said wells are adapted for holding volumes of protein solution from about 0.001 nl to about 250 nl and are in fluid communication through micro-channels with another plurality of wells for holding precipitate solution.

16. The microarray of Claim 15 wherein said wells comprise a material that is minimally water absorbing.

17. The microarray of Claim 15 further comprising an optically clear path from said wells.

18. The microarray of Claim 15 wherein said wells comprise a material that is substantially hydrophobic.

19. The microarray of Claim 15 wherein at least two channels have different dimensions.

20. The microarray of Claim 15 wherein said wells in said microarray further comprises a means for sealing to prevent evaporation from said wells.

21. The method of Claim 1, wherein said micro-channels are about 0.0001 microns to about 2 microns wide and about 0.00005 to about 0.5 microns deep.

22. The method of Claim 1, wherein said micro-channels are about 0.0001 microns to about 0.2 microns wide and about 0.00005 to about 0.01 microns deep.

23. The method of Claim 1, wherein said wells for protein solution hold a volume of about 0.01 nl to about 10 nl.

24. The microarray of Claim 15, wherein said micro-channels are about 0.0001 microns to about 2 microns wide and about 0.00005 to about 0.5 microns deep.


25. The microarray of Claim 15, wherein said micro-channels are about 0.0001 microns to about 0.2 microns wide and about 0.00005 to about 0.01 microns deep.

26. The microarray of Claim 15, wherein said wells for protein solution hold a volume of about 0.01 nl to about 10 nl.

27. A method as defined in Claim 1 substantially as herein described with reference to any example thereof and with or without reference to the accompanying figures.

28. A microarray as defined in Claim 11 substantially as herein described with reference to any example thereof and with or without reference to the accompanying figures.

29. A microarray as defined in Claim 15 substantially as herein described with reference to any example thereof and with or without reference to the accompanying figures.






Figure 1

2/3




Figure 2A




Figure 2B




Figure 2C




Figure 2D




Figure 2E

3/3




Figure 3




Figure 4

END

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.